AGROSTIS LACUNA-VERNALIS (POOIDEAE: POEAE: AGROSTIDINAE), A NEW SPECIES FROM CALIFORNIA

Paul M. Peterson & Robert J. Soreng

Department of Botany National Museum of Natural History Smithsonian Institution Washington, DC 20013-7012, U.S.A. peterson@si.edu;sorengr@si.edu

Dylan Neubauer

David Styer

P.O. Box 444 Moss Landing, CA 95039, U.S.A. david.styer@sbcglobal.net

Randall Morgan

2026 Back Ranch Road Santa Cruz, California 95060, U.S.A. dneubauer1111@gmail.com

157 Market Street Santa Cruz, California 95060

Vern Yadon

1119 Buena Vista Avenue Pacific Grove, California 93950-5537 vly@mbay.net

ABSTRACT

Agrostis lacuna-vernalis P.M. Peterson & Soreng, sp. nov., is described and illustrated. The new species occurs within or on the margins of vernal pools in Monterey County, California and is known only from the former Fort Ord Army Military Base. The new species is morphologically similar to A. blasdalei but differs by having an annual habit, shorter glumes (1.5-2.4 mm versus 1.8-4 mm), shorter lemmas (1.1-1.5 mm versus 1.5-3 mm), and longer paleas (0.4-0.7 mm versus ±0.3 mm).

RESUMEN

Se describe e illustra una nueva especie, Agrostis lacuna-vernalis P.M. Peterson & Soreng, sp. nov. La nueva especie ocurre dentro o en de los márgenes de piscinas vernales en el condado de Monterey, California y se sabe solamente del anterior la Fort Ord Army Military Base. La nueva especie es morfológicamente similar al A. blasdalei pero diferencia teniendo un hábito anual, los glumas más cortos (1.5-2.4 mm contra 1.8-4 mm), los lemas más cortos (1.1-1.5 mm contra 1.5-3 mm) y los paleas más largos (0.4-0.7 mm contra ±0.3 mm).

A small annual species of Agrostis L. had gone undetected on the former Fort Ord Army Military Base, now Fort Ord Public Lands (administered by the Department of Defense and the Department of the Interior, Bureau of Land Management) in Monterey County, California. Ellen Holmes, Nicolas Jensen, Randall Morgan, Dylan Neubauer, David Styer, and Vern Yadon initially collected this enigmatic grass but were unable to name it. Consequently, PMP and RJS were contacted for help, and based on their knowledge, it did not appear to be any previously described species of Agrostis. Since the new taxon is annual in habit, has narrow panicles, and has spikelets with awnless lemmas that are smooth and glabrous and relatively long paleas, there is nothing similar to it in the Flora of North America (Harvey 1993, 2007) and the second edition of the Jepson Manual (Peterson & Harvey in press). The possibility that the plant might be an introduction was considered, but it does not match any species from elsewhere in the world (Clayton et al. 2006). Earlier work on native coastal California Agrostis by Crampton (1967) recognized A. clivicola Crampton with two varieties, both now placed in synonymy of A. densiflora Vasey, and A. blasdalei var. marinensis Crampton, now placed in synonymy of A. blasdalei Hitchc. (Harvey 1993, 2007; Peterson & Harvey in press).

Agrostis is a member of the subtribe Agrostidinae and tribe Poeae in the subfamily Pooideae (Soreng et al. 2007). The Agrostidinae are distributed globally in temperate regions and tropical mountains, include 16 genera in some 550 species worldwide (Clayton & Renvoize 1986), and can be characterized by single-flowered

J. Bot. Res. Inst. Texas 5(2): 421 - 426. 2011

Journal of the Botanical Research Institute of Texas 5(2)

spikelets, glumes exceeding the lemma, lemmas with lateral veins extending to the apex, usually dorsal awns, hyaline paleas, membranous ligules, and caryopses with a punctiform hila and short embryos. The subtribe includes the diverse, ecologically important, and morphologically difficult genera: *Agrostis* (approximately 220 species), *Calamagrostis* Adans. (approximately 270 species), as well as other smaller genera (Saarela et al. 2010). In this paper we describe this unusual annual species as new to science.

Agrostis lacuna-vernalis P.M. Peterson & Soreng, sp. nov. (Fig. 1). TYPE: U.S.A. CALIFORNIA. Monterrey Co.: Fort Ord Army Military Base, Butterfly Valley, 2.8 km SSW of East Garrison and 9 km SW of Salinas, 36°37' 56.2"N, 121°44' 39.2"W, 140 m, 27 Apr-22 May 2010, R. Morgan, D. Styer & D. Neubauer s.n. (HOLOTYPE: US-3621794!).

Ab Agrostis blasdalei plantis annuis, glumis 1.5-2.4 mm longis, lemmatibus 1.1-1.5 mm longis, paleais 0.4-0.7 mm longis, recedit.

Loosely tufted annuals; major roots 0.1-0.3 mm diam.; culms (1.5-)5-30 cm tall, slender (ca. 0.25 mm diam.), erect to decumbent at the base, or slightly geniculate, smooth; internodes 2 or 3 per culm, nodes yellowishbrown, smooth, glabrous. Vegetative shoots intravaginal, prophylls (within 1-3 cm of the base), flattened and papery, 2 keeled, smooth except on the keels. Leaves mostly cauline and basal; sheaths 3/2 to 3/4 as long as the internodes below, open to the base, margins broadly overlapping; ligules of upper leaves 1.2-1.7 mm long, hyaline, margins decurrent, abaxially lightly scabrous, apex acute to obtuse, erose, scabrous; blades 1.4-5(-11) mm long, 0.3-1(-1.5) mm wide, flat, folded to loosely involute on drying, thin and delicate, margins moderately minutely scaberulous, abaxially lightly scaberulous. Panicles 1-7.5 cm long, 1-6 mm wide, narrow, spicate, with 12-90(-110) spikelets, 1-3 branches per node; branches mostly 0.6-1.4 cm long, strict, appressed to suberect, capillary, moderately scabrous, hooks not in distinct lines; pedicels shorter or up to 2.5x longer than the spikelets, distally slightly expanded 0.2-0.3 mm just below the spikelet. Spikelets 1.5-2.4 mm long, 1-flowered; rachilla extension absent; glumes 1.5-2.4 mm long, longer than the floret, subequal (first slightly larger), ovate, herbaceous to scarious, greenish to distinctly purplish, 1-veined, laterally compressed, smooth to minutely muriculate below, scaberulous on the upper sides or only near the apex, more coarsely scaberulous along mid-vein from near the base, the minute hooks whitish under 10-50x magnification, apex mostly acute, scaberulous, margins lightly scabrous with slender hooks from near the middle to the apex; callus distinct, short, blunt, smooth, glabrous, disarticulation scar round; lemmas 1.1-1.5 mm long, subchartaceous, fragile, glossy, obscurely 5-veined, smooth, glabrous, clear to light green with a purplish blush near the vein tips and sometimes on the sides, unawned, keel vein extending to near the apex, the lateral veins faintly visible near the apex where there is commonly some purple coloration, margins broadly overlapping to within 0.15 mm of the base, smooth, apex obtuse to truncate, erose with age; paleas 0.7-1.1 mm long, about 1/2-2/3 as long as the lemma, scarious-hyaline, smooth, obscurely 2 veined, the veins closely spaced, the outer margin broader than the inter-keel gap, apex obtuse and minutely erose; stamens 3, anthers 0.4-0.7 mm long, yellowish; lodicules 2, 0.4-0.5 mm long, hyaline, slightly acentrically s-shaped, elliptic-lanceolate, entire, glabrous; ovary glabrous with two terminal adjacent styles and two stigmas. Caryopses 0.9-1.2 mm long, sub-adherent to the palea, ellipsoid, ventrally longitudinally distinctly grooved, dorsiventrally compressed, light greenish-brownish, translucent, embryo about 1/8 the length of the fruit, hilum indistinct, endosperm slightly malleable (lipid present).

Comments.—The new species can be distinguished from other species of *Agrostis* in North America by the following combination of characters: annual habit, narrow spicate panicles, spikelets with a glabrous callus, unawned lemmas that are smooth and subchartaceous, and anthers 0.4–0.7 mm long. The species is named after the habitat, vernal pool; "lacuna" for pool, and "vernalis" for spring or ephemeral.

Distribution and habitat.—Agrostis lacuna-vernalis is known from Mima mound area in Butterfly Valley (Fort Ord Army Military Base) and Machine Gun Flats (Fort Ord Public Lands) on the Monterey Peninsula between 115–145 meters elevation. It grows within or on the margins of vernal pools and is associated with (*= exotic/introduced): *Aira caryophyllea L., Allium hickmanii Eastw., *Avena barbata Link, *Briza minor L., Brodiaea terrestris Kellogg ssp. terestris, *Bromus hordeaceus L., Calochortus uniflorus Hook. & Arn., Castilleja ambigua ssp. insallutata (Jeps.) T.I. Chuang & Heckard, Cicendia quadrangularis (Lam.) Griseb., *Cotula coroPeterson et al., A new species of Agrostis from California



FIG. 1. Agrostis lacuna-vernalis [R. Morgan, D. Styer & D. Neubauer s.n. (US)]. A. Habit. B. Branch. C. Spikelet. D. Lemma, lateral view. E. Floret. F. Lemma, ventral view. G. Lodicules, stamens, pistil, and palea. H. Palea, dorsal view. I. Lodicules, pistil, and palea. J. Caryopsis, ventral view. K. Caryopsis, lateral view. L. Caryopsis, dorsal view.

nopifolia L., Danthonia californica Bol., Deschampsia danthonioides (Trin.) Benth., Elatine brachysperma A. Gray, Eleocharis acicularis (L.) Roem. & Schult., Eryngium armatum (S. Watson) J.M. Coult. & Rose, Isoetes howellii Engelm., Juncus bufonius (L.) var. bufonius, *J. capitatus Weigel, Lasthenia conjugens Greene, Leptosiphon parviflorum Benth., Lilaea scilloides (Poir.) Hauman, *Lolium multiflorum Lam., *Lythrum hyssopifolia L., Microseris paludosa (Greene) J.T. Howell, Montia fontana L., Plagiobothrys chorisianus var. hickmanii (Greene) I.M. Johnst., Pilularia americana A. Braun, *Plantago coronopus L., P. elongata Pursh, P. erecta E. Morris, Psilocarphus chilensis A. Gray, Ranunculus californicus Benth., Sisyrinchium bellum S. Watson, Trifolium barbigerum Torr., T. buckwestiorum Isely, T. polyodon Greene, T. truncatum (Greene) Greene, T. variegatum Nutt., and *Vulpia bromoides (L.) Gray.

Additional specimens examined. U.S.A. California. Monterey Co.: Butterfly Valley, 24 May 2011, D. Styer, C. Schneider, S. Hubbard, N. Wigington & K. Kasunich 1 (JEPS); D. Styer, C. Schneider, S. Hubbard, N. Wigington & K. Kasunich 2 (CAS); D. Styer, C. Schneider, S. Hubbard, N. Wigington & K. Kasunich 3 (US); D. Styer, C. Schneider, S. Hubbard, N. Wigington & K. Kasunich 4 (RSA); Machine Gun Flats, 27 May 2011, D. Styer s.n. (JEPS, US).

DISCUSSION

The new species can be distinguished from other species of Agrostis in North America and California by possessing the following combination of characters: annual habit, narrow spicate panicles, spikelets with a glabrous callus, unawned lemmas, palea 1/2-2/3 the lemma, and anthers 0.4-0.7 mm long. Agrostis lacuna-vernalis is morphologically similar to three other California natives, A. blasdalei, A. densiflora, and A. variabilis Rydb. (Peterson & Harvey in press; Table 1). The new species differs from A. blasdalei by having an annual habit, shorter glumes (1.5-2.4 mm versus 1.8-4 mm), shorter lemmas (1.1-1.5 mm versus 1.5-3 mm), and longer paleas (0.4–0.7 mm versus ±0.3 mm). It differs from A. densiflora by having an annual habit, narrower blades [0.3-1 (-1.5) mm versus 2-10 mm], shorter lemmas (1.1-1.5 mm versus 1.5-2 mm), and a glabrous callus (versus minutely hairy). It differs from A. variabilis by having an annual habit, shorter lemmas (1.1-1.5 mm versus 1.5-2 mm), well-developed paleas (0.4-0.7 mm versus absent or minute), and a glabrous callus (versus minutely hairy). The new species also resembles Agrostis muelleriana Vickery from Australia and New Zealand but differs by having both cauline and basal leaf blades (mostly basal in A. muelleriana), narrower leaf blades [0.3-1(-1.5) mm versus 1-5 mm], shorter ligules (1.2-1.7 mm versus 2-3 mm), shorter glumes [1.5-2.4 mm versus (2-)2.2-3.5 mm), shorter lemmas 1.1-1.5 mm versus 1.3-2(-3) mm], and longer paleas (0.4-0.7 mm versus vestigial or absent) [Jacobs 2009]. Although we consider the new species to belong to Agrostis s.s., generic concepts in Agrostis s.l. have only been superficially examined by DNA studies (Soreng et al. 2007; Saarela et al. 2010), and their taxonomy can be controversial. The Poaceae editor for the Jepson Manual: Vascular Plants of California, ed. 2 (JMVPC2) treats in Agrostis one annual we prefer to treat as Bromidium tandilense (Kuntze) Rúgolo, and two perennials we prefer to classify as Lachnagrostis filiformis (G. Forst.) Trin. and Podagrostis thurberiana (Hitchc.) Hultén. The separation of these genera from Agrostis was/is accepted in the Catalogue of New World Grasses (Soreng et al. 2003, 2011) and was accepted for the Flora of North America (Barkworth et al. 2007). Under the imposed JMVPC2 generic concept there are now five annual species of Agrostis in California. Below we present a modified key to these annuals, adding the new species to the Agrostis treatment of Peterson & Harvey (in press) in JMVPC2. The new species is globally rare. It is locally frequent, but only known from within a 2.5 km² area, and then only within wetlands in that area. There are estimated to be between 1000 and 10,000 individuals. Its maximum potential range is limited; similar, more or less intact Mima mound/vernal pool habitat covers less than 5 km² in the region (estimated from Google, DitigalGlobe aerial photo projection). Possible threats include competition from exotic grasses and other plants, and destruction of habitat from road building, off road vehicles, domestic livestock grazing, and adverse changes to the drainage patterns.

Peterson et al., A new species of Agrostis from California

TABLE 1. Morphological and habitat comparison of Agrostis lacuna-vernalis to similar California grasses A. blasdalei, A. densiflora, and A. variabilis.

Species	A. blasdalei	A. densiflora	A. variabilis	A. lacuna-vernalis
Longevity	perennial	perennial	short-lived perennial	annual
Lemma length (mm) Lemma awn (mm)	1.5–3 occasionally present, 4–7.5, geniculate	1.5–2 occasionally present, < 3.5, straight	1.5–2 generally absent	1.1-1.5 absent
Lemma texture & vestiture	scareous, easily torn or punctured, mat to sub-lustrous, smooth, nerves distinct	scareous, easily torn or punctured, mat to sub-lustrous, smooth, nerves distinct	scareous, easily torn or punctured, mat to sub-lustrous, smooth,	subchartaceous, glossy, smooth, easily fractured, nerves obscure

Palea length (mm) Palea/Lemma ratio Anther length (mm) Callus

Panicle density Panicle exertion

Longest branches (cm)Elevation (m) Habitat

ca. 0.3 < 1/31-2

glabrous

dense base generally included in upper sheath

< 0.5

< 100 coastal bluffs, dunes, scrub, gravelly soils

0.5 - 0.7 $\pm 1/3$ ±0.5

minutely hairy

dense well exerted

< 0.5

< 200 coastal bluffs, cliffs, scrub, sandy soils

nerves distinct

absent or minute n/a or <1/10 0.5 - 0.7

minutely hairy on the margins fairly dense well exerted

0.5 - 1.5

1600-4000 meadows, talus slopes, subalpine forests, and alpine 0.7 - 1.11/2-2/3 0.4-0.7 glabrous

fairly dense well exerted

0.6 - 1.4

115-145 vernal pools around Mima mounds

KEY TO THE ANNUAL SPECIES OF "AGROSTIS" IN CALIFORNIA

1. Lemmas awnless	A. lacuna-vernalis
1. Lemmas awned from back or near the tip.	
2. Panicles open, generally oblong to ovate in outline; spikelets not crowded, panicle axes clearly visible	A. elliottiana Schult.
2. Panicles dense, generally cylindrical; spikelets crowded, overlapping, panicle axes not clearly visible.	
3. Lemma teeth 4, two < 1 mm long, other two 1-1.5 mm; lemmas awned below middle; callus densely s	hort-hairy
A. tai	ndilensis (Kuntze) Parodi
Lemma teeth 0 or 2, equal in length; lemmas awned at or above middle; callus generally sparsely h minute.	airy, hairs
4. Lemmas 2–4 mm long, awn 8–10 mm long	_ A. hendersonii Hitchc.
4. Lemmas 1.5-2 mm long, awn 3.5-8 mm long	A. microphylla Steud.

ACKNOWLEDGMENTS

We thank Alain Touwaide for help preparing the Latin diagnosis, Alice R. Tangerini for illustrating the new species, and anonymous reviewers for suggesting improvements to the manuscript.

REFERENCES

BARKWORTH, M.E, K.M. CAPELS, S. LONG, L.K. ANDERTON, AND M.B. PIEP. 2007. Flora of North America north of Mexico, volume

24. Oxford University Press, New York and Oxford.

CLAYTON, W.D. AND S.A. RENVOIZE. 1986. Genera Graminum. Grasses of the world. Kew Bull. Add. Ser. 13:1-389. CLAYTON, W.D., M.S. VORONTSOVA, K.T. HARMAN, AND H. WILLIAMSON. 2006 onwards. GrassBase - The online World grass flora. The Board of Trustees, Royal Botanic Gardens, Kew. http://www.kew.org/data/grasses-db.html (accessed 17 Dec 2010).

CRAMPTON, B. 1967. New Agrostis from the California coast. Brittonia 19:174-177.

HARVEY, M.J. 1993. Agrostis. In: J.C. Hickman, ed. The Jepson manual: higher plants of California. University of California Press, Berkeley. Pp. 1227-1231.

HARVEY, M.J. 2007. Agrostis L. In: M.E. Barkworth, K.M. Capels, S. Long, L.K. Anderton, and M.B. Piep, eds. Flora of North

Journal of the Botanical Research Institute of Texas 5(2)

America north of Mexico. Magnoliophyta: Commelinidae (in part): Poaceae, part 1. Oxford University Press, New York and Oxford. 24:633–662.

- JACOBS, S.W.L. 2009. Agrostis. In: A. Wilson, ed. Flora of Australia, volume 44A, Poaceae 2. CSIRO Publishing, Melbourne, Australia. Pp. 163–173.
- PETERSON, P.M. AND M.J. HARVEY. In press. Agrostis. In: B.G. Baldwin, D.H. Goldman, D.J. Keil, R. Patterson, T.J. Rosatti, and D.H. Wilken, eds. The Jepson manual: vascular plants of California, second edition. University of California Press, Berkeley. Pp. 1413–1417.
- SAARELA, J.M., Q. LIU, P.M. PETERSON, R.J. SORENG, AND B. PASZKO. 2010. Phylogenetics of the grass 'Aveneae-type plastid clade' (Poaceae, Pooideae, Poeae) based on plastid and nuclear ribosomal sequence data. In: O. Seberg, G. Petersen, A.S. Barfod, and J.I. Davis, eds. Diversity, phylogeny, and evolution in the Monocotyledons. Aarhus University Press, Descent. Do. 557, 597.

Denmark. Pp. 557–587.

- SORENG, R.J., J.I. DAVIS, AND M.A. VOIONMAA. 2007. A phylogenetic analysis of Poaceae tribe Poeae sensu lato based on morphological characters and sequence data from three plastid-encoded genes: evidence for reticulation, and a new classification for the tribe. Kew Bull. 62:425–454.
- SORENG, R.J., P.M. PETERSON, G. DAVIDSE, E.J. JUDZIEWICZ, F.O. ZULOAGA, T.S. FILGUEIRAS, AND O. MORRONE. 2003. Catalogue of New World grasses (Poaceae): IV. subfamily Pooideae. Contr. U.S. Natl. Herb. 48:1–730.
- SORENG, R.J., P.M. PETERSON, G. DAVIDSE, E.J. JUDZIEWICZ, F.O. ZULOAGA, T.S. FILGUEIRAS, AND O. MORRONE. 2011. Catalogue of New World grasses (Poaceae). http://www.tropicos.org/Project/CNWG (accessed June 2011).

